


Agromet Advisory Service bulletin 05/12 dt. 31-01-12
 (Issued jointly by IAAS Vellanikkara,
 Kerala Agricultural University, India Meteorological Dept. & Agricultural dept.)

THRISSUR

Significant weather for the preceding week at Vellanikkara



Rainfall weekly total	0.0mm
Maximum temperature	31.9-33.2 ⁰ C
Minimum temperature	19.5-21.9 ⁰ C
Max. RH	57-80%
Min . RH	29-36%
Wind speed	6.1-11.1 km/hr
Sun shine hour	8.6-10.2 hour
Evaporation	4.9 – 6.9 millimeter



CURRENT SYNOPTIC CONDITION:

A feeble trough of low pressure lies over Lakshadweep area.

MULTIMODEL ENSEMBLE BASED DISTRICT LEVEL WEATHER FORECAST

ISSUED ON: 31.1.2012

VALID TILL 08:30 HRS IST OF THE NEXT 5 DAYS

STATE : KERALA

PARAMETERS

ENSEMBLE FCST

	DAY-1 01/02	DAY-2 02/02	DAY-3 03/02	DAY-4 04/02	DAY-5 05/02
--	----------------	----------------	----------------	----------------	----------------

DISTRICT: THRISSUR

Rainfall (mm)	0	0	0	0	0
Max Temperature (deg C)	33	33	33	33	33
Min Temperature (deg C)	22	22	22	22	22
Total cloud cover (octa)	3	3	4	4	5
Max Relative Humidity (%)	80	80	81	82	84
Min Relative Humidity (%)	42	44	39	44	45
Wind speed (kmph)	006	006	006	005	005
Wind direction (deg)	50	90	90	90	320
7 days cumulative rainfall for the period from 1/2 to 7/2 - 0					

Crop stage

Puncha paddy is started in certain area. Nendran banana is four months old. Basin mulching is done in coconut, banana, arecanut etc... Harvesting is started in Cool season vegetables.

Agro Meteorological Advisories

Paddy	If Puncha paddy is taken after Mundakan, two weeks time should be given for the land preparation as well as for decaying of the straw which remains in the field. Varieties having short to medium duration are suggested for Puncha e.g. Jyothi, Mattathriveni, Manupriya, Varsha, Gouri etc While preparing the land, organic manure is applied @ 2 tons per one acre.
-------	--

Coconut	To control coried bug, apply 0.1% carbaryl suspension on the on the newly opened inflorescence after the respective phase of the female flowers and spray e entire crown excluding the leaves and older branches.
Cabbage and cauliflower	Drench the soil with trichoderma (20g per litre of water) or psuedamonus (20g per litre of water) or Copper oxychloride (3 g per litre of water) with a interval of 15days to control diseases. Leaf eating insects are controlled by 3% neem oil. Bacterial soft rot of the curd is controlled by spraying psuedamonus (10g per one liter of water) from the curd emergence onwards.
Banana	Mulch banana basins using 3.5 kg of hay or dried materials. To control Sigatoka leaf spot, cut and burn all severely affected leaves. Spray 1% Bordeaux mixer soon after the appearance of the initial symptoms. To control pseudostem weevil remove the dry outer sheaths of the pseudostem of all infected and un-infested plants in the endemic areas and spray any of the recommended insecticides like quinalphos 0.05% or carbaryl 0.2%. Drenching all the leaf axils, rhizome and surrounding soil and all round the entire pseudostem inserting the nozzle through the bore holes made by the larvae if any and also within the outer sheathes
Amaranth	As far as possible, avoid use of insecticides or fungicides. To control rust disease, spray 10g pseudomonas per one liter of water or fresh cow dung extract @ 20g per one liter of water.
Vegetables Land preparation	Apply Neemoil – Garlic emulsion for control of sucking pests. Preparation of Pancha paddy is started in certain areas. Nendran banana is four to five months old. V